

## ERRATUM

JOANNA B. FAWCETT

### Locally triangular graphs and normal quotients of the $n$ -cube

*Example 4.4.* Here I claim that the halved graphs of  $\Pi = (Q_n)_K$  are not isomorphic by Lemma 4.1(i) since  $|\Pi_2(0^K)| = 13$  while  $|\Pi_2(e_1^K)| = 14$ , but this is not a very good proof! However, it is still the case that these graphs are not isomorphic: a tedious calculation shows that the halved graph of  $\Pi$  containing  $0^K$  is regular with valency 13, while the halved graph of  $\Pi$  containing  $e_1^K$  is regular with valency 14.

*Proof of Theorem 1.1.* On the last line of the proof, I claim that the group  $K$  is unique up to conjugacy in  $\text{Aut}(Q_n)$  by Theorem 1.4. This claim is true, but Theorem 1.4 is not enough. I wish to prove the following: if  $\Gamma$  is a halved graph of  $(Q_n)_K$  and  $(Q_n)_L$  for some even  $K, L \leq \text{Aut}(Q_n)$  such that  $d_K \geq 7$  and  $d_L \geq 7$ , then  $K$  and  $L$  are conjugate in  $\text{Aut}(Q_n)$ . By Theorem 1.4, it suffices to show that  $(Q_n)_K \simeq (Q_n)_L$ . Since  $\Gamma$  is connected and locally  $T_n$ , it is a halved graph of a unique bipartite rectagraph by the comment after [1, Lemma 4.1], and since  $(Q_n)_K$  and  $(Q_n)_L$  are both bipartite rectagraphs, it follows that  $(Q_n)_K \simeq (Q_n)_L$ , as desired.